

Circulating Serotonin, Catecholamines, and Central Nervous System Circuitry Related to Some Cardiorespiratory, Vascular, and Hematological Disorders

Fuad Lechin, MD, PhD*
Bertha van der Dijs, MD*
Alex E. Lechin, MD†

**Universidad Central de Venezuela, Institute of Experimental Medicine, Sections of Neurochemistry, Neurophysiology, Neuroimmunology and Neuropharmacology, Faculty of Medicine, Caracas, Venezuela*

†*University of Houston, Department of Clinical Science, Houston, Texas*

KEY WORDS: ANS assessment, arterial hypertension, bronchial asthma, carcinoid syndrome, chronic congestive heart failure, idiopathic thrombocytopenic purpura, platelet disorders, polycythemia vera, pulmonary hypertension

ABSTRACT

This review presents results that show that assessment of circulating neurotransmitters is the most effective and, indeed imperative, way to attain a scientific approach to the autonomic nervous system (ANS). The resulting ANS profile makes it possible to prescribe neuropharmacologic therapies to correct central nervous system circuitry disorders responsible for the abnormalities registered at the peripheral ANS level. Some examples of disastrous therapeutic

trials based on poor knowledge of neuropharmacology and the ANS in general are provided. In addition, data regarding the protagonist role exerted by circulating plasma serotonin in pulmonary bronchoconstriction, pulmonary vasoconstriction, and platelet serotonin in both cardiovascular and hematological disorders are given. All of these disorders were dramatically improved by the administration of neuropharmacologic agents that affected both platelet and neuronal serotonin uptake.

INTRODUCTION

A great deal of published research presents results dealing with the assessment of the autonomic nervous system (ANS). Physiologic, pharmacologic, and clinical parameters have been tested in order to reach a reliable approach to that target. Cardiovascular and/or circu-

lating catecholamines are the most frequently assessed parameters used to evaluate the “so called” peripheral ANS. This over-simplified concept underlies the “two-faces coin” paradigm: sympathetic versus parasympathetic activity. This peripheral neuroautonomic profile does not take into account that the central nervous system (CNS) circuitry, which includes noradrenergic, adrenergic, dopaminergic, serotonergic, and acetylcholinergic neuronal nuclei, displays physiologic interactions that redound in peripheral ANS activity. Hence, the assessment of ANS activity should be carried out on the basis of the measurement of all, and not some, circulating neurotransmitters released from glandular plus neural sources.

In addition to the above, these parameters should be measured during basal as well as after the application of different physiologic stimuli. The rationality of these testing procedures is based on the well-known fact that neural sympathetic, adrenal sympathetic, and parasympathetic branches respond differently to specific stimuli under both physiologic and pathophysiologic situations.¹⁻⁷

This review summarizes the results obtained from the assessment of circulating neurotransmitters as well as other physiologic parameters in 30,000 normal and diseased persons throughout the past 25 years.⁸⁻²³ In addition, the effects of many neuropharmacologic drugs whose CNS and peripheral-acting mechanisms that have been well established are also examined. These procedures assessed the particular contribution of the neural sympathetic, adrenal sympathetic, and parasympathetic systems to the peripheral neuroautonomic profile of the subjects investigated. In addition, the neurophysiologic and neuropharmacologic mechanisms that shed some light on the CNS-peripheral autonomic crosstalk are discussed.

CIRCULATING SEROTONIN

Circulating serotonin (5-hydroxytryptamine or 5-HT) arises from the enterochromaffin cells. Parasympathetic drive plus hormonal- plus food-derived chemicals trigger 5-HT release to both intestinal lumen and portal circulation.^{24,25} In addition, circulating catecholamines are also able to activate enterochromaffin cells.²⁶ The fraction of intestinal 5-HT released to the peripheral circulation is cleared by the liver and lungs.²⁷ Finally, all 5-HT that escapes from the liver plus lungs clearance is taken up and stored by platelets (p-5-HT).²⁸ However, a very small fraction of circulating 5-HT remains free in the plasma (f-5-HT).

f-5-HT is rapidly taken up by the lungs (in vitro perfusion systems), which are able to clear 70% to 90% of exogenously infused 5-HT during one passage through the pulmonary circulation.²⁹ An intact endothelium, particularly in the microvasculature, seems to be a prerequisite for normal clearance of 5-HT, and 5-HT is metabolized within these cells.³⁰ In human airways, 5-HT is localized at nerve endings in the pulmonary neuroendocrine cells (PNEC)³¹ and may be released upon exposure to local airway conditions such as hypoxia, hyperoxia, and hypercapnia.^{32,33} PNEC are granulated epithelial cells and can be detected throughout the lung from the trachea to the alveoli.³⁴ PNEC constitute the vagal presynaptic element and the vagal nerve endings, the postsynaptic element.³⁵

Total circulating serotonin (5-HT) includes platelet-serotonin (p-5-HT; normal values (NV) = 150-250 ng/mL) plus free-serotonin in the plasma (f-5-HT; NV= 0.7-1.5 ng/mL). Both p-5-HT and f-5-HT increase during postprandial periods⁸ and other hyperparasympathetic (physiologic or pathophysiologic) situations.³⁶⁻³⁸ Increased release from enterochromaffin cells accounts for the

p-5-HT rise,³⁷⁻³⁹ whereas increased levels of circulating acetylcholine (ACh) that interfere with platelet uptake of 5-HT are responsible for augmented f-5-HT plasma levels.^{1,2} Conversely, increased circulating catecholamines are responsible for the lowered p-5-HT and the increased f-5-HT registered during stressful situations that trigger platelet aggregation.^{2,38}

CIRCULATING CATECHOLAMINES

Two different pools constitute circulating catecholamines: adrenal gland secretions and neural sympathetic release. Adrenal gland catecholamines consist of adrenaline (Ad) (75% to 80%), plus noradrenaline (NA) and dopamine (DA), which account for the remaining 20% to 25%. However, some fraction of circulating catecholamines can be taken up by sympathetic terminals, to be further released together with the NA plus DA synthesized in these terminals.⁴⁰ Neural sympathetic release of catecholamines comprises 80% to 90% NA plus 10% to 20% DA. Both NA and DA are synthesized in separate pools. The latter is released before the former and exerts a modulatory role by acting on DA₂ receptors existing at sympathetic terminals.⁴¹⁻⁴³

CNS CIRCUITRY RESPONSIBLE FOR SEROTONERGIC PLUS CATECHOLAMINERGIC ACTIVITIES

Although circulating 5-HT does not cross the blood brain barrier, a close and complex anatomic plus physiologic crosstalk takes place between central and peripheral 5-HT activities, as well as among serotonergic plus catecholaminergic and parasympathetic activities. However, this review refers only to some well-established findings considered basic to a general understanding by readers.

The CNS-5-HT system includes six neuronal nuclei located at the midline

throughout the pontine-midbrain and medullary regions: the dorsal raphe (DR), median raphe (MR), periaqueductal gray (PAG), raphe magnus (RM), raphe obscurus (RO), and raphe pallidus (RP).^{44,45} These serotonergic nuclei display direct and indirect modulation on all sympathetic and parasympathetic (medullary, hypothalamic, and spinal) nuclei.⁴⁶ This is possible due to complex polysynaptic mechanisms that modulate the peripheral ANS and depend on a broad web of direct and indirectly driven information (crosstalk). For instance, the nucleus RO sends direct excitatory axons to the medullary motor vagal complex (dorsal motor + nucleus ambiguus).⁴⁷⁻⁵⁰ In turn, the 5-HT-RO receives excitatory sensory vagal nuclei (dorsal + area postrema).⁵¹ The area postrema is located outside the blood brain barrier and thus is accessible to the blood stream. This nucleus is crowded with 5-HT₃ receptors that can be excited by plasma 5-HT.⁵² According to this, any f-5-HT increase in the plasma triggers parasympathetic activation, which in turn, will increase ACh plasma levels.^{53,54} Considering that plasma ACh interferes with 5-HT uptake by platelets, any f-5-HT increase in the plasma will trigger further excitation of the area postrema. This excitatory drive is transmitted to the motor vagal complex and vagal nerves, whose discharge at intestinal enterochromaffin cells triggers 5-HT release. This parasympathetic positive feedback will result in a further increase of circulating ACh, which in turn, will boost f-5-HT plasma levels.^{53,54} This positive serotonergic-parasympathetic feedback constitutes the called Bezold-Jarisch reflex, responsible for uncontrollable symptoms such as nausea, vomiting, diarrhea, blood pressure decrease, etc.

Other examples of the central-peripheral crosstalk may be seen. For instance, PAG and MR serotonergic

nuclei send excitatory and inhibitory axons, respectively, to the C1-Ad (adrenaline neurons) located at the rostral ventrolateral medullary areas.⁵⁵⁻⁵⁹ These C1-Ad nuclei are directly responsible for adrenal gland secretion as shown by retrovirus tracing.⁶⁰ Conversely, a RP serotonergic nucleus sends excitatory axons to spinal sympathetic preganglionic neurons located at the intermediolateral columns of the thoracic plus lumbar sympathetic segments. These preganglionic (ACh) sympathetic neurons send ACh-preganglionic axons to the postganglionic (NA) cells, located at the sympathetic ganglia, whose axons constitute the sympathetic nerves.^{61,62} Thus, peripheral sympathetic activity is made up of two separate systems, ruled by two different CNS circuits.

In short, the sympathetic preganglionic neurons that send axons to adrenal glands are directly ruled by C1-Ad medullary nuclei, while the sympathetic preganglionic neurons sending axons to sympathetic ganglia belong to a separate circuit and receive direct axons from the NA(A5) plus NA(A6) pontine nucleus.⁶³ However, both central preganglionic sympathetic neurons receive serotonergic modulatory axons. According to the above, two different branches of peripheral sympathetic activities exist, and these two branches converge to form the pool of plasma catecholamines.⁶⁴⁻⁶⁹ One part of that pool arises from NA-DA released from sympathetic nerves, while the other pool (Ad + DA + NA) arises from adrenal gland secretion.⁶⁵⁻⁷¹

PULMONARY VASOCONSTRICTION

Pulmonary vasoconstriction is the main pathophysiological disorder in both primary and secondary pulmonary hypertension. It is characterized by raised pulmonary vascular resistance, which results in diminished right-heart function due to increased right ventricular afterload. Pathogenic mechanisms that

deal with this syndrome focus on abnormalities in the interaction between endothelial and smooth muscle cells that require mediators favoring vasoconstriction. Pulmonary artery branches are compliant structures with few muscle fibers, allowing the pulmonary vascular bed to function as a high-flow, low-pressure circuit. Many circulating vasoactive mediators have been postulated to play a part in pulmonary hypertension, including 5-HT,⁷²⁻⁷⁶ which has been found raised during acute periods. With respect to this, plasma catecholamines and indolamines (5-HT) in 11 primary pulmonary hypertension cases plus two cases associated with scleroderma and three associated with Raynaud disease have been investigated. In addition, 16 patients with secondary pulmonary hypertension (two with vasculitis, three with chronic bronchitis, nine with chronic bronchial asthma, and two with obesity) have been examined. All of these patients were investigated during relapse as well as relief periods. Although the supine resting plus orthostasis plus exercise test was performed during the latter period, only supine-resting assessment was performed during relapse.⁷⁷⁻⁸¹

The results showed that although all patients with pulmonary hypertension showed raised f-5-HT plasma values during relapses, opposite profiles were registered during relief periods. Patients who presented with scleroderma and Raynaud disease had very high NA/Ad plasma ratio and very low plasma tryptophane levels. Conversely, the other patients (those with vasculitis, chronic bronchitis, obesity, and bronchial asthma) with pulmonary hypertension but not scleroderma or Raynaud disease showed low NA/Ad ratio (<2) and normal tryptophane plasma levels. These opposing plasma neurotransmitter profiles showed an absolute neurosympathetic over adrenal sympathetic

predominance in the former whereas the opposite profile is seen in the latter group. These two plasma catecholamine profiles are similar to those found in patients with essential and nonessential hypertension, respectively.^{9,10,71}

The fact that a small dose of oral tianeptine (12.5 mg), a drug that enhances 5-HT uptake, suppressed acute symptoms in patients with pulmonary hypertension within the first hour of its administration showed that the f-5-HT plasma peak was responsible for worsening the symptoms. It is logical to assume that the abrupt f-5-HT rise registered in secondary pulmonary hypertension is triggered by increased platelet aggregability, secondary to increased plasma levels of Ad that were registered in these patients. However, this factor is absent in patients with primary pulmonary hypertension. The fact that these patients showed greatly raised plasma levels of DA during pulmonary hypertension relapses should be noted, in accordance with the incontrovertible fact that not only ACh, but also DA, interferes with platelet serotonin uptake.^{82,83} Obviously, the circulating DA rise that registered in patients with primary pulmonary hypertension during attacks came from the DA pool existing at sympathetic terminals, which is not only co-secreted with NA but is released before NA during neural sympathetic discharge.^{42,43} For instance, it is known that NA(A6) plus NA(A5), DR-5-HT, and MR-5-HT pontine neuronal nuclei show progressive reduction of their “firing activity” throughout the four slow wave sleep (SWS) stages and, further, they show zero firing activity during REM sleep.^{84,85} This slope is paralleled by the NA but not the Ad plasma slope.^{11,86,87} NA plasma levels fall progressively and reach minimal levels at the REM sleep stage whereas Ad plasma levels fall abruptly during the first 10 minutes of supine resting position, when

the subjects are still awake. These findings are consistent with the observation that the NA pontine activity fades progressively throughout sleep periods. Patients affected by essential hypertension showed no NA plasma fall during sleep and, moreover, presented no REM sleep stage, which is consistent with the hypernoradrenergic activity found in this type of hypertensive patient.^{9-11,71,86,87}

The investigation of circulating neurotransmitters throughout wake-sleep periods in patients showing the “uncoping” stress profile revealed that they had a very low NA/Ad plasma ratio during wake and sleep stages alike. The NA plasma level fell abruptly during the supine-resting period. They did not present the normal SWS fading and abruptly reached the REM sleep stage.^{12-16,88,89} This is explained by the exhaustion of NA pontine neurons registered in “uncoping” stressed mammals.⁸⁸ Plasma catecholamine values are greater than normal during REM sleep stage in “uncoping” stressed mammals; however, the NA/Ad ratio shows minimal values at this period because of the absolute absence of neural sympathetic activity. It is consistent with the knowledge that nocturnal cardiovascular, cerebrovascular, and respiratory attacks usually occur at REM periods in stressed subjects who show maximal C1-Ad over the NA pontine neurons predominance at this sleep stage.

The f-5-HT/p-5-HT plasma ratio registered during REM sleep is an appropriate parameter of the parasympathetic activity when tested in normal subjects. This f-5-HT/p-5-HT ratio profile is paralleled by the low blood pressure and minimal pulse rate registered in normal subjects. These findings are consistent with the fact that both NA and Ad reach minimal levels at this period.¹¹

ACh interferes with platelet uptake, which is consistent with the raised f-5-HT/p-5-HT ratio found during the REM

sleep period, at which time maximal parasympathetic activity is registered. This peripheral parasympathetic hyperactivity would be positively associated with findings showing that the midbrain and medullary ACh neurons display maximal activity during the REM period. However, the raised f-5-HT plasma levels registered throughout the abnormal sleep stages in “uncoping” stressed subjects do not reflect parasympathetic activity but platelet aggregation, secondary to the high Ad plasma levels they always present. These findings should be viewed as examples of the CNS-peripheral autonomic crosstalk.

An additional concern is expressed here regarding the identification of CNS-autonomic disorders triggered by the consumption of sleeping pills and antidepressants that provoke a distortion of the wake-sleep cycle and, consequently, trigger physiologic disorders in both CNS and peripheral autonomic systems.⁹⁰⁻⁹³

ARTERIAL HYPERTENSION

Two types of hypertensive subjects, according to their plasma neurotransmitter profiles were found: patients with essential hypertension and those with nonessential hypertension. The former group showed a greater than normal NA/Ad plasma ratio and low tryptophane plasma levels, whereas the latter showed a low NA/Ad ratio (<2) and normal or raised tryptophane plasma levels. f-5-HT and plasma DA were normal in the former group, while f-5-HT and plasma Ad were elevated in the second group. A close positive correlation was found between DA versus f-5-HT in patients with essential hypertension, whereas a close positive correlation was also found between Ad versus f-5-HT in the nonessential hypertension group.^{9,10,71} Because DA interferes with p-5-HT uptake^{42,43,94} and Ad triggers platelet aggregation, this agrees with the above-

mentioned correlations, respectively.³

Because the rat model of essential hypertension presents increased activity of NA pontine neurons⁹⁵⁻¹⁰⁰ and the rat model of “uncoping” stress situation is characterized by exhausted NA pontine neurons plus hyperactive C1-Ad nuclei,^{4,101} these two hypertensive profiles might be extrapolated to humans.^{9,10,71,100,101}

CHRONIC CONGESTIVE HEART FAILURE

Although patients with chronic congestive heart failure (CCHF) show raised catecholamine circulating levels, their plasma NA/Ad ratio is very low, revealing absent NA pontine activity. The failure by Swedberg et al,¹⁰² who attempted to treat these patients with an imidazoline agonist on the grounds that patients with CCHF presented “so called” hyper-sympathetic activity, occurred because the investigators ignored that the two peripheral sympathetic activities do not constitute a unit, but rather two well-differentiated sympathetic systems that display associated or dissociated activities.^{69,71,103-106}

The appropriate parameter for assessing the two branches of the sympathetic system should be based on the assay of plasma catecholamines in order to find the NA/Ad ratio, and not the sum of NA and Ad. This assessment should be carried out throughout the complete supine resting plus 1-minute orthostasis plus 5-minute moderate exercise test.^{5,69,107} This test has been performed in 30,000 healthy and diseased persons throughout the last 20 years.⁹⁷ The normal value of NA/Ad ratio at 1-minute orthostasis was found to be 3 to 4. The diastolic blood pressure, but not the systolic blood pressure, should peak in normal subjects at orthostasis. This diastolic blood pressure peak parallels NA plasma rise and directly depends on the NA pontine neuronal firing, which

activates lumbar sympathetic preganglionic neurons and is transmitted to sympathetic nerves.^{108,109} The 1-minute orthostasis test reflects the normal dissociation between the two branches of sympathetic activities.^{5,40,69,99,106,109-113} This NA/Ad ratio increases in patients who show NA pontine over C1-Ad predominance, such as neurogenic hypertension,^{9,10,98,99} Raynaud disease,¹¹⁴ scleroderma,¹¹⁴ endogenous depression,^{17,18} primary pulmonary hypertension,^{76,115} and most autoimmune diseases that are labeled as presenting with a TH-1 immunological profile.¹¹⁶ Conversely, the low NA/Ad plasma ratio is registered in diseases presenting an “uncoping” stress neurochemical profile, eg, bronchial asthma, secondary pulmonary hypertension, gastroduodenal ulcer, most types of cancer and infectious diseases, and autoimmune diseases that show a TH-2 cytokine profile.^{6,116,117}

The two types of sympathetic preganglionic neurons are modulated by serotonergic axons, and this is a clear example of serotonergic sympathetic crosstalk. This phenomenon is evident during sleep, during which different stages are paralleled by NA, Ad, f-5-HT, p-5-HT, and tryptophane blood changes. Considering that the neurophysiologic changes of the different NA, Ad, 5-HT, and ACh CNS nuclei that occur throughout the four slow-wave and REM sleep periods are well known, it is possible to investigate the parallels between CNS and circulating neurotransmitters changes.^{6,117-121}

These findings are consistent with the postulated pathogenic role assigned to f-5-HT as the trigger of pulmonary hypertension relapses. These comments are also consistent with the worsening of all types of pulmonary hypertension when treated with drugs that augment f-5-HT plasma levels, like fenfluramine, sertraline, paroxetine, fluoxetine, and other 5-HT uptake-inhibitors.¹²²⁻¹²⁴

Conversely, the dramatic suppression of pulmonary hypertension symptoms induced by tianeptine, a drug that enhances platelet uptake and triggers an abrupt disappearance of f-5-HT from the plasma, gives additional support.^{46,77-79,81}

CIRCULATING 5-HT AND BRONCHOCONSTRICTION

Studies have shown that levels of f-5-HT in plasma are elevated in symptomatic asthmatic patients during attacks.^{15,118-122} In addition, the plasma concentration of f-5-HT in symptomatic patients with asthma correlates positively with clinical status and negatively with pulmonary function (FEV_1).¹²⁵⁻¹²⁹ 5-HT facilitates cholinergic contractions in human airways in vitro through stimulation of both prejunctional 5-HT₃ and 5-HT₄ receptors.¹³⁰⁻¹³⁵

Although symptomatic patients with asthma have increased plasma levels of NA, Ad, DA, f-5-HT, and cortisol when compared with asymptomatic patients with asthma, f-5-HT was the only factor closely associated with clinical severity and pulmonary malfunction. Other data show that f-5-HT is actively transported by the pulmonary endothelial cells, where it is metabolized by the monoamine oxidase enzyme.¹³² In addition, 5-HT causes constriction of both central and peripheral airways when given to vagotomized cats and other mammals. Further data show that inhaled 5-HT induces an acute fall in lung function (>20% in FEV_1) in asthmatic patients but not in normal subjects.^{133,135}

Potential sources of 5-HT within the pulmonary system include platelets and nerve terminal (PNEC) cells. In humans, 5-HT is concentrated in platelets and is released when platelets aggregate. Platelet aggregation occurs during stressful situations and is common in immunological diseases.¹³⁶⁻¹³⁹

Bronchoconstriction evoked by f-5-

HT involves vagal afferent nerves, and it is inhibited by atropine. f-5-HT can induce bronchoconstriction by affecting presynaptic neuronal 5-HT₃ receptors at the medullary area postrema, located outside the blood brain barrier.¹⁴⁰⁻¹⁴⁹ Increased parasympathetic activity is associated with asthma attacks. Parasympathetic activity releases 5-HT at the intestinal level and provokes an increase of blood 5-HT. In addition, hyperparasympathetic activity (as occurs during sleep and postprandial periods) interferes with p-5-HT uptake, which enhances f-5-HT in the plasma, thereby triggering asthma attacks. Buspirone, a drug that increases central and peripheral parasympathetic activity, provokes asthma attacks in patients with asthma.¹⁹ Tianeptine, a drug that enhances 5-HT uptake by platelets and reduces f-5-HT in the plasma, suppresses asthma attacks within the first hour after administration.^{20,21} It is difficult to know if the dramatic suppression of asthma attacks induced by tianeptine should be attributed to peripheral and/or central effects. This drug enhances the uptake of 5-HT by serotonergic terminals. These 5-HT axons release 5-HT at the medullary respiratory center responsible for the activation of this function.¹⁵⁰⁻¹⁵²

CIRCULATING 5-HT AND PLATELET DISORDERS

Refractory Idiopathic

Thrombocytopenic Purpura

Refractory idiopathic thrombocytopenic purpura (ITP) is a severe pathophysiologic disorder characterized by the development of autoantibodies that destroy platelets. It is included among the TH-2 autoimmune diseases in accordance with the cytokine profiles registered in the blood of these patients.¹⁵³⁻¹⁵⁷ Patients with ITP who do not improve with immunosuppressant therapy are considered refractory, and they usually undergo splenectomy. Nevertheless, an

important percentage of splenectomized patients with ITP show severe and uncontrollable relapses. The assessment of circulating catecholamines and indolamines performed at the Institute of Experimental Medicine, Universidad Central de Venezuela, on those refractory patients with ITP revealed that there was no correlation between the number of platelets and the amount of p-5-HT. In addition, the bleeding disorders correlated negatively with the p-5-HT values rather than with the platelet count.¹⁵⁷⁻¹⁶⁰ Finally, all of these patients showed increased levels of f-5-HT as well as very high levels of plasma Ad, which was responsible for their very low NA/Ad plasma ratio. Because the administration of tianeptine, a drug that enhances 5-HT uptake by platelets, triggered a dramatic disappearance of bleeding episodes, this supports the hypothesis that 5-HT plays a primordial role in platelet activation. However, a definite improvement of patients with ITP was achieved by the normalization of the NA/Ad ratio with neuropharmacologic therapy.¹⁵⁷

Polycythemia Vera

Polycythemia vera (PV) is a myeloproliferative disorder of unknown cause. PV is associated with a NA over 5-HT predominance that is paralleled by a TH-1 over TH-2 immune profile;¹¹⁶ this led the investigators to prescribe neuropharmacologic therapy to reverse this neuroautonomic unbalance.¹⁶¹ Normalization of the neurochemical CNS disorder triggered by neuropharmacologic therapy was followed by restoration of both the normal ANS profile and the immunological peripheral balance.

It is important to emphasize the role played by p-5-HT as a thrombogenic factor. It has been shown that the platelet content of 5-HT rather than the platelet count itself is the most important factor involved in hemosta-

sis.^{157,162,163} The disappearance of thrombotic episodes in a patient with PV occurred 3 weeks after start of neuropharmacologic therapy, at which time p-5-HT, but not the platelet count, dropped.^{116,161} This finding is the mirror image of that observed in purpura thrombocytopenic patients, in whom the bleeding stoppage was paralleled by the p-5-HT increase rather than the platelet number increase.^{157,162-164} This phenomenon was discussed in several papers, and it assigned a fundamental role to p-5-HT in all types of thrombotic events, which can be eliminated by lowering platelet 5-HT content with drugs that interfere with 5-HT uptake by both platelets and serotonergic neurons.^{158-160,165} Finally, reversion of the TH-1 immunologic profile seen in patients with PV following reversion of the CNS-NA over 5-HT predominance suggests that this immunologic disorder underlies the etiopathogenesis of this disease.

The spleen bone marrow and all hematopoietic organs receive neural sympathetic innervation. Sympathetic nerves arise from the lumbar spinal sympathetic neurons that send preganglionic axons to sympathetic ganglia. Such preganglionic branches contact postganglionic sympathetic neurons (located at this level), which innervate the aforementioned hematopoietic organs. In turn, spinal sympathetic preganglionic neurons receive axons from the NA pontine nuclei, which is the main center responsible for neural sympathetic activity. These facts explain the beneficial therapeutic effects exerted by CNS-acting drugs that potentiate serotonergic neurons that bridle NA pontine nuclei in improving patients with PV.

Carcinoid Syndrome

Carcinoid syndrome is another example of adrenal over neural sympathetic predominance plus raised levels of circulating p-5-HT and f-5-HT, as well as a

TH-2 immunological profile. These patients were treated with neuropharmacologic therapy to revert the neuroimmunologic profiles.^{166,167} Both neuroautonomic and immunologic disorders should be included among the pathophysiologic mechanisms underlying this disease (TH-2 profile). This inference is reinforced by all clinical, ANS, and immunologic improvements that were triggered by neuropharmacologic manipulations similar to those used to revert Ad over NA predominance.

The raised circulating 5-HT level found in carcinoid patients agrees with the enterochromaffin cell overactivity that is always present in this disease. Furthermore, because p-5-HT remained elevated whereas the f-5-HT was normalized in these patients after clinical improvement, and although their enterochromaffin cells remained overactive, the symptoms appear to have been related to the raised f-5-HT plasma levels. Both the physiologic disorders and their clinical symptoms were paralleled by adrenal over neural sympathetic predominance periods. This phenomenon may be associated with the maximal ability of the neural sympathetic system to annul the parasympathetic drive, which is responsible for enterochromaffin cell secretion and for increased f-5-HT levels, all of which occurs during excessive parasympathetic drive. The increase of f-5-HT registered in these circumstances may be exacerbated because this indolamine excites 5-HT₃ and 5-HT₄ receptors located at the medullary area postrema (outside the blood brain barrier), which is connected to the motor vagal complex.¹⁰⁰ This results in further increase of the peripheral parasympathetic drive discharge over the enterochromaffin cells (Bezold-Jarisch reflex). Such mechanisms may explain the hyperserotonergic storm that occurs in carcinoid patients.

The enhancement of natural killer

cells (NK-cell) cytotoxicity as well as the shifting of TH-2 to TH-1 immunologic profile, triggered by the neuropharmacologic therapy prescribed to these carcinoid patients, merits special mention. In 1987, neuropharmacologic therapy was able to improve patients with different types of cancer.²² In addition, the clinical improvement was paralleled by an increase in NK cell activity.^{40,45,168,169} These reports were further confirmed in other studies.^{16,170-171} It was also shown that clinical severity in patients with cancer correlated negatively with NA/Ad ratio, whereas clinical improvement showed positive correlation with NA/Ad. The foregoing findings are in line with those by other investigators who showed that some drugs which improved colon and lung cancer triggered NK cell activation.¹⁷⁰

CONCLUSION

Extensive experimental, clinical, and therapeutic data have been presented that show that the assessment of all (but not one or two) circulating neurotransmitters is the only effective way to accurately evaluate the peripheral ANS. Furthermore, a reliable evaluation depends on assessing neurotransmitters not only during basal (supine-resting) but also at orthostasis and exercise stress periods. Data demonstrating successful therapeutical approaches to bronchial asthma, pulmonary vasoconstriction, carcinoid syndrome, and platelet disorders were presented. In addition, data dealing with the peripheral and CNS autonomic disorders that underlie the pathophysiology of CCHF and arterial hypertension were shown. Furthermore, broad scientific data dealing with the CNS circuitry responsible for controlling the peripheral ANS, depending on a close crosstalk between them were offered. Finally, the administration of drugs that act at the CNS level should be managed by physicians who possess a thorough knowledge of CNS

physiology. These results are supported not only by some recent therapeutic trial failures but also by the ongoing iatrogenic neuropharmacologic medication prescribed by doctors seeking to treat depression, sleep disorders, and other psychiatric disorders, whose ANS repercussions are often ignored.

REFERENCES

1. Rausch JL, Janowsky DS, Risch SC, Huey LY. Physostigmin effects on serotonin uptake in human blood platelets. *Eur J Pharmacol.* 1985;109:91-96.
2. Vesifeld IL, Vasiljiev VN, Iljicheva RF. Relationship of catecholamines, histamine and serotonin in men under different kinds of stress. In: Usdin E, Kvetnansky R, Kopin IJ, eds. *Catecholamines and Stress.* Oxford, UK: Pergamon Press; 1976:527-528.
3. Larsson PT, Hjemdhal P, Olsson G, et al. Altered platelet function during mental stress and adrenaline infusion in humans: evidence for an increase aggregability in vivo as measured by filtragemetry. *Clin Sci.* 1989;76:369-376.
4. Breier A, Albus M, Pickar D, et al. Controllable and uncontrollable stress in humans: alterations in mood and neuroendocrine and psychophysiological function. *Am J Psychiatry.* 1987;144:1419-1425.
5. Young JB, Rowe JW, Pallota JA, et al. Enhanced plasma norepinephrine response to upright posture and oral glucose administration in elderly human subjects. *Metabolism.* 1980;29:532-535.
6. Irwin M, Thompson J, Miller C, et al. Effects of sleep and sleep deprivation on catecholamines and interleukin-2 levels in humans: clinical implications. *J Clin Endocrinol Metab.* 1999;84:1979-1985.
7. Robertson D, Johnson GA, Robertson RM, et al. Comparative assessment of stimuli that release neuronal and adrenomedullary catecholamines in man. *Circulation.* 1979;59:637-643.
8. Lechin F, van der Dijs B, Lechin M, et al. Effects of an oral glucose load on plasma neurotransmitters in humans: involvement of REM sleep? *Neuropsychobiology.* 1992;26:4-11.
9. Lechin F, van der Dijs B, Lechin ME, et al. Plasma neurotransmitters throughout oral glucose tolerance test in non-depressed essential hypertension patients. *Clin Exp Hypertens.* 1993;15:209-240.

10. Lechin F, van der Dijs B, Lechin ME, et al. Plasma neurotransmitters, blood pressure and heart rate during supine-resting, orthostasis, and moderate exercise conditions in two types of hypertensive patients. *Res Comm Biol Psychol Psychiatry*. 1997;22:97-112.
11. Lechin F, Pardey-Maldonado B, van der Dijs B, et al. Circulating neurotransmitters during the different wake-sleep stages in normal subjects. *Psychoneuroendocrinology*. 2004;29:669-685.
12. Lechin F, van der Dijs B, Rada I, et al. Plasma neurotransmitters and cortisol in duodenal ulcer patients. Role of stress. *Dig Dis Sci*. 1990;35:1313-1319.
13. Lechin F, van der Dijs B, Jackubowitz D, et al. Effects of clonidine on blood pressure, noradrenaline, cortisol, growth hormone and prolactin plasma levels in low and high intestinal tone subjects. *Neuroendocrinology*. 1985;40:253-261.
14. Lechin F, van der Dijs B, Orozco B, et al. Plasma neurotransmitters, blood pressure and heart during supine-resting, orthostasis and moderate exercise in severely ill patients: a model of failing to cope with stress. *Psychother Psychosom*. 1996;65:129-136.
15. Lechin AE, Varon J, van der Dijs B, Lechin F. Plasma catecholamines and indoleamines during attacks and remission on severe bronchial asthma: possible role of stress. *Am J Respir Crit Care Med*. 1994;149:778.
16. Lechin F, van der Dijs B, Vitelli G, et al. Psychoneuroendocrinological and immunological parameters in cancer patients: involvement of stress and depression. *Psychoneuroendocrinology*. 1990;15:435-451.
17. Lechin F, van der Dijs B, Orozco B, et al. Plasma neurotransmitters, blood pressure and heart rate during supine-resting, orthostasis and moderate exercise conditions in major depressed patients. *Biol Psychiatry*. 1995;38:166-173.
18. Lechin F, van der Dijs B, Jakubowicz D, et al. Effects of clonidine on blood pressure, noradrenaline, cortisol, growth hormone, and prolactin plasma levels in high and low intestinal tone depressed patients. *Neuroendocrinology*. 1985;41:156-162.
19. Lechin F, van der Dijs B, Jara H, et al. Effects of buspirone on plasma neurotransmitters in healthy subjects. *J Neural Transm*. 1998;105:561-573.
20. Lechin F, van der Dijs B, Lechin A, et al. The serotonin uptake-enhancing drug tianeptine suppresses asthmatic symptoms in children. A double-blind crossover placebo-controlled study. *J Clin Pharmacol*. 1998;38:918-925.
21. Lechin F, van der Dijs B, Orozco B, et al. Neuropharmacological treatment of bronchial asthma with an antidepressant drug: tianeptine. A double-blind crossover placebo-controlled study. *Clin Pharmacol Ther*. 1998;64:223-232.
22. Lechin F, van der Dijs B, Azocar J, et al. Stress, immunology and cancer: effect of psychoactive drugs. *Arch Ven Farm Clin Terap*. 1987;6:28-43.
23. Lechin F, van der Dijs B, Lechin A, et al. Doxepin therapy for postprandial symptomatic hypoglycaemic patients: neurochemical, hormonal and metabolic disturbances. *Clin Sci (Lond)*. 1991;80:373-384.
24. Ahlman H, Bhargava HN, Dahlstrom A, et al. On the presence of serotonin in the gut lumen and possible release mechanisms. *Acta Physiol Scand*. 1981;112:263-269.
25. Burks TF, Long JP. Catecholamine-induced release of 5-hydroxytryptamine from perfused vasculature of isolated dog intestine. *J Pharm Sci*. 1966;55:1383-1386.
26. Larsson I, Ahlman H, Bhargava HN, et al. The effects of splanchnic nerve stimulation on the plasma levels of 5-HT and SP in the portal vein of the cat. *J Neural Transm*. 1979;46:105-112.
27. Kjellstrom T, Ahlman H, Dahlstrom A, Risberg B. The cellular uptake of 5-hydroxytryptamine in the rat lung in vivo—a histochemical study. *Acta Physiol Scand*. 1982;116:455-460.
28. Born GVR, Gillson RE. Studies on the uptake of 5-hydroxytryptamine by blood platelets. *J Physiol (Lond)*. 1959;146:472-491.
29. Davis RB, Wang Y. Rapid pulmonary removal of 5-hydroxytryptamine in the intact dog. *Proc Soc Exp Biol*. 1965;118:799-800.
30. Gillis CN, Cronau LH, Mandel S, Hammond GL. Indicator dilution measurements of 5-hydroxytryptamine clearance by human lung. *J Appl Physiol*. 1979;46:1178-1183.
31. Lauweryns JM, van Ranst L, Lloyd RV, O'Connor DT. Chromogranin in bronchopulmonary neuroendocrine cells. Immunocytochemical detection in human, monkey, and pig respiratory mucosa. *J Histochem Cytochem*. 1987;35:113-118.
32. Keith IM, Will JA. Dynamics of the neuroendocrine cell—regulatory peptide system in the lung. Specific overview and new results. *Exp Lung Res*. 1982;3:387-402.
33. Moosav H, Smith P, Heath D. The Feyrter cell in hypoxia. *Thorax*. 1973;28:729-741.
34. Johnson DE, Georgieff MK. Pulmonary neuroendocrine cells. Their secretory products and their potential roles in health and chron-

- ic lung disease in infancy. *Am Rev Respir Dis.* 1989;140:1807-1812.
35. Levitt RC, Mitzner W. Autosomal recessive inheritance of airway hyperreactivity to 5-hydroxytryptamine. *J Appl Physiol.* 1989;67:1125-1132.
 36. Tobe T, Izumikawa F, Sano M, Tanak C. Release mechanisms of 5-HT in mammalian gastrointestinal tract—especially vagal release of 5-HT. In: Fujita T, ed. *Endocrine Gut-Pancreas.* Amsterdam, Netherlands: Elsevier; 1976:371-380.
 37. Schworer H, Racke K, Kilbinger H. Cholinergic modulation of the release of 5-hydroxytryptamine from the guinea pig ileum. *Naunyn Schmiedebergs Arch Pharmacol.* 1987;336:127-132.
 38. Lechin F, van der Dijs B, Lechin ME. Plasma neurotransmitters and functional illness. *Psychother Psychosom.* 1996;65:293-318.
 39. Gronstad KO, Zinner MJ, Dahlstrom A. Vagal release of serotonin into gut lumen and portal circulation via separate control mechanisms. *J Surg Res.* 1987;43:205-210.
 40. Lechin F, van der Dijs B, Amat J, Lechin ME. Neuroanatomical basis: the noradrenergic system. In: Lechin F, van der Dijs B, eds. *Neurochemistry and Clinical Disorders: Circuitry of Some Psychiatric and Psychosomatic Syndromes.* Boca Raton, FL: CRC Press; 1989:2-9.
 41. Lokhandwala MF, Hedge SS. Cardiovascular dopamine receptors: role of renal dopamine and dopamine receptors in sodium excretion. *Pharmacol Toxicol.* 1990;66:237-243.
 42. Mannelli M, Pupilli C, Fabbri G, et al. Endogenous dopamine (DA) and DA₂ receptors: a mechanism limiting excessive sympathetic-adrenal discharge in humans. *J Clin Endocrinol Metab.* 1988;66:626-631.
 43. Mercurio G, Gessa G, Rivano CA, et al. Evidence for a dopaminergic control of sympathoadrenal catecholamines release. *Am J Cardiol.* 1988;62:827-828.
 44. Lechin F, van der Dijs B, Lechin E. Pharmacology of the autonomic nervous system imbalances. In: Lechin F, van der Dijs B, Lechin E, eds. *The Autonomic Nervous System: Physiological Basis of Psychosomatic Therapy.* Barcelona, Spain: Editorial Cientifico-Medica; 1979:119-150.
 45. Lechin F, van der Dijs B, Amat J, Lechin ME. Neuroanatomical basis: the serotonergic system. In: Lechin F, van der Dijs B, eds. *Neurochemistry and Clinical Disorders: Circuitry of Some Psychiatric and Psychosomatic Syndromes.* Boca Raton, FL: CRC Press; 1989:10-12.
 46. Lechin F, van der Dijs B, Lechin ME: Bronchial asthma and pulmonary hypertension. In: Lechin F, van der Dijs B, Lechin ME, eds. *Neurocircuitry and Neuroautonomic Disorders. Reviews and Therapeutic Strategies.* Basel, Switzerland: Karger; 2002:66-68.
 47. Feldman PD. Electrophysiological effects of serotonin in the solitary tract nucleus of the rat. *Arch Pharmacol.* 1994;349:447-454.
 48. McCann M, Hermann GE, Rogers RC. Nucleus raphe obscurus (nRO) influence vagal control of gastric motility in rats. *Brain Res.* 1989;486:181-184.
 49. Jacobs BL, Fornal CA. Activity of brain serotonergic neurons in the behaving animal. *Pharmacol Rev.* 1991;43:563-578.
 50. Sawchenko PE. Central connections of the sensory and motor nuclei of the vagus nerve. *J Auton Nerv Syst.* 1983;9:13-26.
 51. Thurston CL, Helton ES. Effects of intravenous phenylephrine on blood pressure, nociception, and neural activity in the rostral ventral medulla in rats. *Brain Res.* 1996;717:81-90.
 52. Miller AD, Nonaka S. Mechanisms of vomiting induced by the serotonin-3 receptor agonists in the cat: effect of vagotomy, splanchnicectomy or area postrema lesion. *J Pharmacol Exp Ther.* 1992;260:509-517.
 53. Callera JC, Bonagamba LGH, Sevoz C, et al. Cardiovascular effects of microinjection of low doses of serotonin into the NTS of unanesthetized rats. *Am J Physiol.* 1997;272:R1135-R1142.
 54. Ireland SJ, Tyers MB. Pharmacological characterization of 5-hydroxy-tryptamine-induced depolarization of the rat isolated vagus nerve. *Br J Pharmacol.* 1987;90:229-238.
 55. Vertes RP, Fortin WJ, Crane AM. Projections of the median raphe nucleus in the rat. *J Comp Neurol.* 1999;407:555-582.
 56. Thurston-Stanfield CL, Ranieri JT, Vallabhapurapu R, Barnes-Noble D. Role of vagal afferents and the rostral ventral medulla in intravenous serotonin-induced changes in nociception and arterial blood pressure. *Physiol Behav.* 1999;57:753-767.
 57. Yusof APM, Coote JH. Patterns of activity in sympathetic postganglionic nerves to skeletal muscle, skin and kidney during stimulation of the medullary raphe area of the rat. *J Auton Nerv Syst.* 1988;24:71-79.
 58. Farkas E, Jansen ASP, Loewy AD. Periaqueductal gray matter projection to vagal preganglionic neurons and the nucleus tractus solitarius. *Brain Res.* 1997;764:257-261.

59. Larsen PD, Zhong S, Gebber GL, Barman SM. Differential pattern of spinal sympathetic outflow in response to stimulation of the caudal medullary raphe. *Am J Physiol Regul Integr Comp Physiol.* 2000;279:R210-R221.
60. Li YW, Wesselingh SL, Blessing WW. Projections from rabbit caudal medulla to C1 and A5 sympathetic premotor neurons, demonstrated with phaseolus leucoagglutinin and herpes simplex virus. *J Comp Neurol.* 1992;317:379-395.
61. Behbehani MM. Functional characteristics of the midbrain periaqueductal gray. *Prog Neurobiol.* 1995;46:575-605.
62. Chen S, Aston-Jones G. Anatomical evidence for inputs to ventrolateral medullary catecholaminergic neurons from the midbrain periaqueductal gray of the rat. *Neurosci Lett.* 1995;195:140-144.
63. Gauthier P. Pressor responses and adrenomedullary catecholamines release during brain stimulation in the rat. *Can J Physiol Pharmacol.* 1981;59:485-492.
64. Benarroch EE, Balda MS, Finkielman S, Nahmod VE. Neurogenic hypertension after depletion of norepinephrine in anterior hypothalamus induced by 6-hydroxydopamine administration into the ventral pons: role of serotonin. *Neuropharmacology.* 1981;22:29-34.
65. Dampney RAL, Czachurski J, Dembowski K, et al. Afferent connections and spinal projections of the pressor region in the rostral ventrolateral medulla of the cat. *J Auton Nerv Syst.* 1987;20:73-86.
66. Privitera PJ, Granata AR, Underwood MD, et al. C1 area of the rostral ventrolateral medulla as a site for the central hypotensive action of propranolol. *J Pharmacol Exp Ther.* 1988;246:529-533.
67. McCall RB, Harris LT. 5-HT₂ receptor agonists increase spontaneous sympathetic nerve discharge. *Eur J Pharmacol.* 1988;151:113-116.
68. Funakoshi K, Kadota T, Atobe Y, et al. Serotonin-immunoreactive axons in the cell column of sympathetic preganglionic neurons in the spinal cord of the filefish *Stephanolepis cirrhifer*. *Neurosci Lett.* 2000;280:115-118.
69. Young JB, Rosa RM, Landsberg L. Dissociation of sympathetic nervous system and adrenal medullary responses. *Am J Physiol.* 1984;247:E35-E40.
70. Astier B, Kitahama K, Denoroy L, et al. Biochemical evidence for an interaction between adrenaline and noradrenaline neurons in the rat brainstem. *Brain Res.* 1986;397:333-340.
71. Lechin F, van der Dijs B, Lechin AE. Neural sympathetic activity in essential hypertension. *Hypertension.* 2004;44:e3-e4.
72. Hervé P, Launay JM, Scrobohaci ML, et al. Increased plasma serotonin in primary pulmonary hypertension. *Am J Med.* 1995;99:249-254.
73. Deuchar GA, Docherty A, MacLean MR, Hicks MN. Pulmonary hypertension secondary to left ventricular dysfunction: the role of nitric oxide and endothelin-1 in the control of pulmonary vascular tone. *Br J Pharmacol.* 2002;135:1060-1068.
74. Belohlávková S, Simák J, Kokesová A, et al. Fenfluramine-induced pulmonary vasoconstriction: role of serotonin receptors and potassium channels. *J Appl Physiol.* 2001;91:755-761.
75. Strange JW, Wharton J, Phillips PG, Wilkins MR. Recent insights into the pathogenesis and therapeutics of pulmonary hypertension. *Clin Sci.* 2002;102:253-268.
76. Humbert M, Labrune P, Sitbon O, et al. Pulmonary arterial hypertension and type-1 glycogen-storage disease: the serotonin hypothesis. *Eur Respir J.* 2002;20:59-65.
77. Lechin F, van der Dijs B, Lechin AE. Pulmonary hypertension, left ventricular dysfunction and plasma serotonin. *Br J Pharmacol.* 2002;137:937-938.
78. Lechin F, van der Dijs B. Serotonin and pulmonary vasoconstriction. *J Appl Physiol.* 2002;92:1363-1364.
79. Lechin F, van der Dijs B, Lechin AE. Plasma serotonin, pulmonary hypertension and bronchial asthma. *Clin Sci.* 2002;103:345-346.
80. Lechin F, van der Dijs B. Plasma serotonin, bronchial asthma and tianeptine. *Clin Physiol.* 2001;21:723-724.
81. Lechin F, Lechin AE, van der Dijs B. Tianeptine, plasma serotonin, and pulmonary hypertension. *Lancet.* 2003;361:87-88.
82. Gordon JL, Olverman JH. 5-Hydroxytryptamine and dopamine transport by rat and human blood platelets. *Br J Clin Pharmacol.* 1978;62:219-226.
83. Lingjaerde O, Kildemo O. Dopamine uptake in platelets: two different low-affinity, saturable mechanisms. *Agents Actions.* 1981;11:410-416.
84. McGinty DJ, Harper RM. Dorsal raphe neurons, depression of firing during sleep in cats. *Brain Res.* 1976;101:569-575.
85. MacCarley RW. REM sleep and depression, common neurobiological control mechanisms. *Am J Psychiat.* 1982;139:5-13.

86. Lechin F, van der Dijs B: Slow wave sleep (SWS), REM sleep (REMS), and depression. *Res Commun Psychol Psychiatr Behav.* 1984;9:227-262.
87. Lechin F, van der Dijs B, Lechin ME. Central neurocircuitry functioning during the wake-sleep cycle. In: Lechin F, van der Dijs B, Lechin ME, eds. *Neurocircuitry and Neuroautonomic Disorders. Reviews and Therapeutic Strategies.* Basel, Switzerland: Karger; 2002:3-13.
88. Glavin GB. Selective noradrenaline depletion markedly alters stress responses in rats. *Life Sci.* 1985;37:461-465.
89. Lechin F, van der Dijs B, Benaim M. Stress versus depression. *Prog Neuropsychopharmacol Biol Psychiatry.* 1996;20:899-950.
90. Richelson E. Pharmacology of antidepressants—characteristics of the ideal drug. *Mayo Clin Proc.* 1994;69:1069-1081.
91. Grignaschi G, Invernizzi RW, Fanelli E, et al. Citalopram-induced hypophagia is enhanced by blockade of 5-HT_{2c} receptors. *Br J Pharmacol.* 1998;124:1781-1787.
92. Ko H-C, Lu R-B, Shiah I-S, Hwang C-C. Plasma free 3-methoxy-4-hydrophenylglycol predicts response to fluoxetine. *Biol Psychiatry.* 1997;41:774-781.
93. Lechin F, van der Dijs B, Benaim M. Benzodiazepines: tolerability in elderly patients. *Psychother Psychosom.* 1996;65:171-182.
94. Rabey JM, Oberman Z, Graff E, Korczyn AD. Decreased dopamine uptake into platelet storage granules in Gilles de la Tourette disease. *Biol Psychiatry.* 1995;38:112-115.
95. Elam M, Thoren P, Svensson TH. Locus coeruleus neurons and sympathetic nerves: activation by visceral afferents. *Brain Res.* 1986;375:117-125.
96. Tsukamoto K, Sved AF, Ito S, et al. Enhanced serotonin-mediated responses in the nucleus tractus solitarius of spontaneously hypertensive rats. *Brain Res.* 2000;863:1-8.
97. Koulu M, Saavedra JM, Niwa M, Linnoila M. Increased catecholamines metabolism in the locus coeruleus of young spontaneously hypertensive rats. *Brain Res.* 1986;369:361-364.
98. Mancia G, Grassi G, Parati G, Zanchetti A. The sympathetic nervous system in human hypertension. *Acta Physiol Scand.* 1997;640(suppl):117-121.
99. Lambert GW, Kaye DM, Thompson JM, et al. Internal jugular venous spillover of noradrenaline and metabolites and their association with sympathetic nervous activity. *Acta Physiol Scand.* 1998;163:155-163.
100. Lechin F, van der Dijs B, Villa S, Lechin AE. Central nervous system (CNS) circuitry involved in blood pressure regulation. In: Lechin F, van der Dijs B, eds. *Neurochemistry and Clinical Disorders: Circuitry of Some Psychiatric and Psychosomatic Syndromes.* Boca Raton, FL: CRC Press; 1989:121-150.
101. Burchfield SR. The stress response: a new perspective. *Psychosom Med.* 1979;41:661-672.
102. Swedberg K, Brislow M, Cohn JN, et al. Effects of sustained-release moxonidine, an imidazoline agonist, on plasma norepinephrine in patients with chronic heart failure. *Circulation.* 2002;105:1797-1803.
103. Abdallah H, Guyenet PG. Prototypical imidazoline-1 receptor ligand monoxidine activates alpha-2 adrenoceptors in bulbo spinal neurons of the RVL. *J Neurophysiol.* 2000;83:766-776.
104. Wu M-F, Gulyani SA, Yau E, et al. Locus coeruleus neurons: cessation of activity during cataplexy. *Neurosci.* 1999;91:1389-1399.
105. Lechin F, Lechin ME, van der Dijs B. Plasma catecholamines and chronic congestive heart failure. *Circulation.* 2002;106:222.
106. Gillette RN, Punnen S, Krieger AJ, Sapru HN. Interdependence of rostral and caudal ventrolateral medullary areas in the control of blood pressure. *Brain Res.* 1984;321:169-174.
107. Lechin F, van der Dijs B, Lechin ME. Plasma neurotransmitters and functional illness. *Psychother Psychosom.* 1996;65:293-318.
108. Yazawa M, Ikeda SI, Ushiyama M, Yanagisawa N. Noradrenergic nerve fibers of the rectal mucosa in autonomic disorders: comparison of histochemical study with clinical severity and changes in plasma noradrenaline induced by standing. *J Neurol Sci.* 1991;104:222-229.
109. Page ME, Valentino RJ. Locus coeruleus activation by physiological challenges. *Brain Res Bull.* 1994;35:557-560.
110. Cedarbaum JM, Aghajanian GK. Noradrenergic neurons of the locus coeruleus: inhibition by epinephrine and activation by the alpha-antagonist piperoxane. *Brain Res.* 1976;112:413-419.
111. Engberg G, Elam M, Svensson TH. Effect of adrenaline synthesis inhibition on brain noradrenaline neurons in locus coeruleus. *Brain Res.* 1981;223:49-58.
112. Hokfelt T, Johansson O, Goldstein M. Central catecholamines neurons as revealed by immunohistochemistry with special reference to the adrenaline neurons. In: Bjorklund A, Hokfelt T, eds. *Handbook of Neuroanatomy, Classical Transmitters in the CNS.* New York, NY: Elsevier Science; 1984;2:157-276.

113. Petty MA, Reid JL. Changes in noradrenaline concentration in brain stem and hypothalamic nuclei during the development of renovascular hypertension. *Brain Res.* 1977;136:376-380.
114. Pancera P, Sansone S, Presciuttini B, et al. Autonomic nervous system dysfunction in sclerodermic and primary Raynaud's phenomenon. *Clin Sci.* 1999;96:49-57.
115. Hoeper MM, Galie N, Simonneau G, Rubin LJ. New treatments for pulmonary arterial hypertension. *Am J Respir Crit Care Med.* 2002;165:1209-1216.
116. Lechin F, van der Dijs B, Lechin ME. Neuroautonomic, neuroendocrine and neuroimmune interactions. In: Lechin F, van der Dijs B, Lechin ME, eds. *Neurocircuitry and Neuroautonomic Disorders. Reviews and Therapeutic Strategies.* Basel, Switzerland; Karger; 2002:57-59.
117. Jarnicki AG, Fallon PG. T helper type-2 cytokine responses: potential therapeutic agents. *Curr Opin Pharmacol.* 2003;3:449-455.
118. Culebras A. The neurology of sleep. *Neurology.* 1992;42(suppl 6):6-8.
119. Ramage AG. Central cardiovascular regulation and 5-hydroxytryptamine receptors. *Brain Res Bull.* 2001;56:425-439.
120. Senba E, Matsunaga K, Tohyama M, Noguchi K. Stress-induced c-fos expression in the rat brain: activation mechanism of sympathetic pathway. *Brain Res Bull.* 1993;31:329-344.
121. Nosjean A, Callera JC, Bonagamba L, et al. Serotonin receptor stimulation in the nucleus tractus solitarius activates non-catecholaminergic neurons in the rat ventrolateral medulla. *Neuroscience.* 2002;112:935-949.
122. Eddahibi S, Adnot S. Anorexigen-induced pulmonary hypertension and the serotonin (5-HT) hypothesis: lessons for the future in pathogenesis. *Respir Res.* 2002;3:9.
123. Lesch KP. Molecular biology, and genetics of the serotonin transporter: psychobiological and clinical implications in serotonergic neurons and 5-HT receptors in the CNS. In: Baumgarten HG, Gothert M, eds. *Handbook of Experimental Pharmacology.* Berlin, Germany: Springer; 1997.
124. Eddahibi S, Raffestin B, Pham I, et al. Treatment with 5-HT potentiates development of pulmonary hypertension in chronically hypoxic rats. *Am J Physiol.* 1997;272:H1173-H1181.
125. Lechin F, van der Dijs B, Orozco B, et al. Increased levels of free serotonin in plasma of symptomatic asthmatic patients. *Ann Allergy Asthma Immunol.* 1996;77:245-253.
126. Lechin F. Asthma, asthma medication and autonomic nervous system dysfunction. *Clin Physiol.* 2001;21:723-724.
127. Lechin F, van der Dijs B, Lechin AE. Severe asthma and plasma serotonin. *Allergy.* 2002;57:258-259.
128. Lechin F. Central and plasma 5-HT, vagal tone and airways. *Trends Pharmacol Sci.* 2000;21:425-426.
129. Lechin F, van der Dijs B, Lechin AE. Tianeptine: a new exploratory therapy for asthma. *Chest.* 2004;125:348-349.
130. Dupont LJ, Pype JL, Demedts MG, et al. The effects of 5-HT on cholinergic contraction in human airways in vitro. *Eur Respir J.* 1999;14:642-649.
131. Cazzola M, Matera MG. 5-HT modifiers as a potential treatment of asthma. *Trends Pharmacol Sci.* 2000;21:13-15.
132. Strum JM, Junod AF. Radioautographic demonstration of 5-hydroxytryptamine-3H uptake by pulmonary endothelial cells. *J Cell Biol.* 1972;54:456-467.
133. Pype JL, Verleden GM, Demedts MG. 5-HT modulates noncholinergic contraction in guinea pig airways in vitro by prejunctional 5/HT1/like receptor. *J Appl Physiol.* 1994;77:1135-1141.
134. Spannake EW, Levin JL, Mellion BT, et al. Reversal 5-HT-induced bronchoconstriction by PG12: distribution of central and peripheral actions. *J Appl Physiol.* 1980;49:521-527.
135. Cushley MJ, Wee LH, Holgate ST. The effect of inhaled 5-hydroxytryptamine (5-HT, serotonin) on airway caliber in man. *Br J Clin Pharmacol.* 1986;22:487-490.
136. Capron A, Joseph M, Ameisen JC, et al. Platelets as effectors in immune and hypersensitivity reactions. *Int Arch Allergy Appl Immunol.* 1987;82:307-312.
137. Cosgriff TM. Viruses and hemostasis. *Rev Infect Dis.* 1989;11(suppl 4):S672-S688.
138. Hechtman HB, Lonergan EA, Staunton HP, et al. Pulmonary entrapment of platelets during acute respiratory failure. *Surgery.* 1978;83:277-283.
139. Cazzola M, Matera MG, Rossi F. Bronchial hyperresponsiveness and bacterial respiratory infections. *Clin Ther.* 1991;13:157-171.
140. Dampney RA. The subretrofacial vasomotor nucleus: anatomical, chemical and pharmacological properties and role in cardiovascular regulation. *Prog Neurobiol.* 1994;42:197-227.
141. Hilaire G, Bou C, Monteau R. Rostral ventrolateral medulla and respiratory rhythmogenesis in mice. *Neurosci Lett.* 1997;224:13-16.

142. Lindsey BG, Arata A, Morris KF, et al. Medullary raphe neurons and baroreceptor modulation of the respiratory motor pattern in the cat. *J Physiol.* 1998;512:863-882.
143. Richard CA, Stremel RW. Involvement of the raphe in the respiratory effect of gigantocellular area activation. *Brain Res Bull.* 1990;25:19-23.
144. Huang Z-G, Subramanian SH, Balnave RJ, et al. Role of periaqueductal grey and nucleus tractus solitarius in cardiorespiratory function in the rat brainstem. *Respir Physiol.* 2000;120:185-195.
145. Holtman JR Jr, Dick TE, Berger AJ. Serotonin-mediated excitation of recurrent laryngeal and phrenic motoneurons evoked by stimulation of the raphe obscurus. *Brain Res.* 1987;417:12-20.
146. Wang Y, Ramage AG. The role of central 5-HT_{1A} receptors in the control of B-fibre cardiac and bronchoconstrictor vagal preganglionic neurons in anesthetized cats. *J Physiol.* 2001;536:753-767.
147. Arita H, Ochiishi M. Opposite effect of 5-hydroxytryptamine on two types of medullary inspiratory neuron with distinct firing patterns. *J Neurophysiol.* 1991;66:285-292.
148. Skaburskis M, Shardonofsky F, Milic-Emili J. Effect of serotonin on expiratory pulmonary resistance in cats. *J Appl Physiol.* 1990;68:2419-2425.
149. Behan M, Brownfield MS. Age-related changes in serotonin in the hypoglossal nucleus of rat: implications for sleep-disordered breathing. *Neurosci Lett.* 1999;267:133-136.
150. Broqua P, Baudrie V, Laude D, Chaouloff F. Influence of the novel antidepressant tianeptine on neurochemical, neuroendocrinological, and behavioral effects of stress in rats. *Biol Psychiatry.* 1992;31:391-400.
151. Piñeyro G, Deveault L, Blier P, et al. Effect of acute and prolonged tianeptine administration on the 5-HT transporter: electrophysiological, biochemical and radiological binding studies in the rat brain. *Naunyn-Schmiedeberg's Arch Pharmacol.* 1995;351:111-118.
152. Lechin F, van der Dijs B, Lechin AE. Treatment of bronchial asthma with tianeptine. *Methods Find Exp Clin Pharmacol.* 2004;26:697-701.
153. Singh VK, Mehrotra S, Agarwal SS. The paradigm of Th1 and Th2 cytokines. Its relevance to autoimmunity and allergy. *Immunol Res.* 1999;20:147-161.
154. Salzet M, Vieau D, Day R. Crosstalk between nervous and immune systems through the animal kingdom: focus on opioid. *Trends Neurosci.* 2000;23:550-555.
155. Downing JEG, Miyan JA. Neural immunoregulation: emerging roles for nerves in immune homeostasis and disease. *Immunol Today.* 2000;21:281-289.
156. Lechin F, van der Dijs B, Lechin ME. Stress. In: Lechin F, van der Dijs B, Lechin ME, eds. *Neurocircuitry and Neuroautonomic Disorders. Reviews and Therapeutic Strategies.* Basel, Switzerland: Karger; 2002:18-22.
157. Lechin F, van der Dijs B, Orozco B, et al. Neuropharmacological treatment of refractory idiopathic thrombocytopenic purpura: roles of circulating catecholamines and serotonin. *Thromb Haemost.* 2004;91:1254-1256.
158. Lechin F, van der Dijs B. Platelet serotonin and thrombostasis. *J Clin Invest.* 2004; Available at <http://www.jci.org> – Letters. April 21.
159. Lechin F, van der Dijs B, Orozco B, et al. Elective stenting, platelet serotonin and thrombotic events. *Platelets.* 2004;15:462-463.
160. Lechin F, van der Dijs B. Platelet aggregation, platelet serotonin and pancreatitis. *JOP.* 2004;5:8001-8003.
161. Lechin F, van der Dijs B, Orozco B, et al. Neuropharmacological therapy of polycythemia vera: roles of circulating catecholamines and serotonin. *Thromb Haemost.* 2005;93:175-177.
162. Gobbi G, Mirandola P, Tazzari PL, et al. Flow cytometry detection of serotonin content and release in resting and activated platelets. *Br J Haematol.* 2003;121:892-896.
163. Dale GI, Friese P, Batar P, et al. Stimulated platelets use serotonin to enhance their retention of procoagulant proteins on the cell surface. *Science.* 2002;415:175-179.
164. Dominguez V, Govezensky T, Gevorkian G, Larralde C. Low platelet counts do not cause bleeding in an experimental model of immune thrombocytopenic purpura in mice. *Haematologica.* 2003;88:679-687.
165. Lechin F, van der Dijs B. Platelet activation and catheter intracoronary brachytherapy [online journal]. *Heart.* 2004; Available at <http://heart.bmjournals.com> Read eLetters) July 1.
166. Lechin F, van der Dijs B, Orozco B, et al. Neuropharmacological therapy of the neuroendocrine carcinoid syndrome: report of two cases. *J Appl Res.* 2005;5:109-114.
167. Lechin F, van der Dijs B. Neuropharmacological therapy of carcinoid syndrome. *Neuroendocrinology.* 2005;81:137-138.
168. Lechin F, van der Dijs B, Lechin S, et al. Neurochemical, hormonal and immunological views of stress: clinical and therapeutic implications in Crohn's disease and cancer.

- In: Velazco M, ed. *Recent Advances in Pharmacology and Therapeutics*. International Congress Series vol. 839. Amsterdam, Netherlands: Excerpta Medica; 1989:57-70.
169. Lechin F, van der Dijs B, Lechin ME. Illustrations of some therapeutic results. In: Lechin F, van der Dijs B, Lechin ME, eds. *Neurocircuitry and Neuroautonomic Disorders. Reviews and Therapeutic Strategies*. Basel, Switzerland: Karger; 2002:75-94.
170. Krause SW, Gastpar R, Andreessen R, et al. Treatment of colon and lung cancer patients with ex vivo heat shock protein 70-peptide-activated, autologous natural killer cells: a clinical phase I trial. *Clin Cancer Res*. 2004;10:3699-3707.
171. Lechin F, van der Dijs B, Lechin AE. Natural killer cells activity and neuroimmunological treatment of cancer. *Clin Cancer Res*. 2004;10:8120-8121.